



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

But the integrand of the first of these two integrals is less than (or equal to) $k^n \sqrt{1 + n^2 k^{n-2}}$ and since the limit of this is 0, the limit of the integral is zero also. To handle the second integral, write the integrand as

$$nx^{2n-1} \sqrt{1 + \frac{1}{n^2 x^{2n-2}}}.$$

We may now develop in series and show that the limit of every term after the first is zero, or we may proceed as follows:

$$\begin{aligned} \int_k^1 nx^{2n-1} \sqrt{1 + \frac{1}{n^2 x^{2n-2}}} dx &= \int_k^1 nx^{2n-1} \left\{ 1 + \sqrt{1 + \frac{1}{n^2 x^{2n-2}}} - 1 \right\} dx \\ &= \int_k^1 nx^{2n-1} \cdot dx + \int_k^1 n \cdot x^{2n-1} \cdot \frac{\frac{1}{n^2 x^{2n-2}}}{\sqrt{1 + \frac{1}{n^2 x^{2n-2}}} + 1} \cdot dx, \end{aligned}$$

by rationalizing the numerator. The limit of the first integral is $\frac{1}{2}$ and since the integrand in the second may be made as small as we please, the limit of the second integral is zero.

Also solved by J. A. CAPARO and the PROPOSER.

379. Proposed by C. N. SCHMALL, New York City.

Express the equation of the folium, $x^3 + y^3 = 3axy$, in parametric form and find the area of the loop.

(From E. B. Wilson's *Advanced Calculus*, p. 296, ex. 5.)

SOLUTION BY E. B. WILSON, Mass. Institute of Technology.

Let $y = mx$, then

$$x = \frac{3am}{1 + m^3}, \quad y = \frac{3am^2}{1 + m^3};$$

the loop being described by values of m from 0 to ∞ . By the formulas for area as a curvilinear integral

$$A = - \int_{m=0}^{\infty} y dx = - \int_0^{\infty} 9a^2 \frac{m^2(1 - 2m^3)dm}{(1 + m^3)^3} = - \int_0^{\infty} 3a^2 \frac{1 - 2u}{(1 + u)^3} du,$$

where $u = m^3$. Then

$$A = - 3a^2 \left[\frac{2}{1 + u} - \frac{3}{2} \frac{1}{(1 + u)^2} \right]_0^{\infty} = \frac{3}{2} a^2.$$

Also solved by ELIJAH SWIFT, C. E. HORNE, WILSON L. MISER, W. C. EELLS, HORACE OLSON, J. A. CAPARO, H. L. AGARD, L. G. WELD, and the PROPOSER.

MECHANICS.

297. Proposed by C. N. SCHMALL, New York City.

A shrapnel shell strikes the ground and then explodes, dispersing its fragments in all directions with a common velocity v . If a be the area of the ground covered by the fragments, and if the dimensions of the shell be neglected, show that $a = \pi v^4/g^2$.

SOLUTION BY HORACE OLSON, Chicago, Illinois.

According to the laws of physics, the range of a projectile on the horizontal plane from which it is thrown is $(2v^2 \sin \theta \cos \theta)/g$ or $(v^2 \sin 2\theta)/g$, θ being the inclination with the horizontal of the line of projection. This range has a maximum value, v^2/g , when θ is 45° . Therefore the area of the ground covered by the fragments is $\pi v^4/g^2$, the area of a circle of radius v^2/g .

Also solved by A. M. HARDING, J. L. RILEY, and P. PEÑALVER.